

Three deformation functors for associative algebras

Frank Schuhmacher

February 2, 2008

Abstract

We show that three deformation functors (deformations of the product, flat deformations and deformations of the relations) assigned to an associative algebra are naturally isomorphic.

Introduction

In the literature (see [1] and references here-in), deformations of an associative algebra A mostly refer to deformations of the associative multiplication of A . But, dually to deformations of schemes and singularities, it is as natural to consider flat deformations of associative algebras. A deformation of the multiplication of A defines directly a flat deformation. In this paper we show that, up to equivalence, any flat deformation over an Artin base is just a deformation of the multiplication.

A third approach is to consider A as a quotient of a free algebra, modulo an ideal of relations, and to study deformations of those relations. To make this approach fit into the flat concept, we also have to control “higher relations” which leads to deforming the differential of a whole free DG algebra resolution of A . By abuse of language, we still speak of deformations of the relations of A . It was shown by Kontsevich [3] that deformations of the associative multiplications of A , parametrized by an Artin algebra B (or by a projective limit of Artin algebras), correspond one-to-one to deformations of the relations of A , parametrized by B . The aim of this paper is to show that flat deformations over Artin algebras are just a third perspective to “deformations of A ”.

Conventions: For the whole paper, fix a ground field k of characteristic zero. We will study deformations of an associative non-unitary k -algebra A . All constructions and statements are true with slight modifications for unitary algebras. The multiplication $A \otimes_k A \longrightarrow A$ will be denoted by α . Deformations will be parametrized by a commutative local artinian algebra $B = (B, \mathfrak{m})$ such that $B/\mathfrak{m} = k$. Let $\mathfrak{Art}$ be the category of these algebras. For any k -module (or DG module) M , set $M_B := M \otimes_k B$ and $M_{\mathfrak{m}} := M \otimes_k \mathfrak{m}$.

1 Deformation theory via DG Lie algebras

A good introduction to deformation theory via DG Lie algebras is given in [5]. We will just recall the general basic concept: To study deformations of a given object A in some category, we assign a deformation functor Def to A , i.e. the set-valued functor $\text{Def} : \mathfrak{Art} \longrightarrow \mathbf{set}$ on the category of Artin algebras, assigning to $B \in \text{Ob}(\mathfrak{Art})$ the quotient of the set of all deformations of A parametrized by B , modulo equivalent deformations. We say that the deformations of A are **governed by** a DG Lie algebra $L = (L, d, [\cdot, \cdot])$, if the functor Def is of the form

$$\text{Def}(B) = \mathcal{MC}(B) / \sim, \quad (1.1)$$

where the right hand-side is defined as follows: The tensor products $L_B := L \otimes B$ and $L_{\mathfrak{m}} := L \otimes \mathfrak{m}$ are again DG Lie algebras with structure maps defined by $d(u \otimes b) = du \otimes b$ and $[u \otimes b, u' \otimes b'] = [u, u'] \otimes bb'$. Since the ideal $\mathfrak{m}$ is nilpotent, the Lie algebra $L_{\mathfrak{m}}^0$ is also nilpotent. We can thus form the group $G := \exp(L_{\mathfrak{m}}^0)$, and the exponential map $\exp : L_{\mathfrak{m}}^0 \longrightarrow G$ is bijective. The group G operates on $L_{\mathfrak{m}}^1$ via

$$\exp(f) \cdot u = \exp(\text{ad}_f)(u),$$

where ad_f is the endomorphism $u \mapsto [f, u]$ of $L_{\mathfrak{m}}^1$. It is convenient to introduce the DGL algebra $\tilde{L}_{\mathfrak{m}} = (\tilde{L}_{\mathfrak{m}}, d_d, [\cdot, \cdot]_d)$, defined by $\tilde{L}_{\mathfrak{m}}^i := L_{\mathfrak{m}}^i$, for $i \neq 1$ and $\tilde{L}_{\mathfrak{m}}^1 := L_{\mathfrak{m}}^1 \oplus k \cdot d$. The structure maps of $\tilde{L}_{\mathfrak{m}}$ are defined by $d_d(u + \lambda d) = du$, $[d, u]_d = du$ and $[d, d]_d = 0$, for $\lambda \in k$ and $u \in L_{\mathfrak{m}}$. We have the affine map $L_{\mathfrak{m}} \longrightarrow \tilde{L}_{\mathfrak{m}}$, given by $u \mapsto \tilde{u} := u + d$. Since $\tilde{L}_{\mathfrak{m}}^0 = L_{\mathfrak{m}}^0$, the group G also acts on $\tilde{L}_{\mathfrak{m}}^1$. Now we can define $\mathcal{MC}(B)$ which is the set

$$\{u \in L_{\mathfrak{m}}^1 \mid du + \frac{1}{2}[u, u] = 0\}$$

of **Maurer-Cartan-elements** of $L_{\mathfrak{m}}$. The tilde-map restricts to a bijection between $\mathcal{MC}(B)$ and the set

$$\tilde{\mathcal{MC}}(B) := \{\tilde{u} \in \tilde{L}_{\mathfrak{m}}^1 \mid [\tilde{u}, \tilde{u}]_d = 0\}.$$

The set $\tilde{\mathcal{MC}}(B)$ is invariant under the action of G . We call two Maurer-Cartan elements $u, v \in \mathcal{MC}(B)$ **(gauge-)equivalent**, if $\tilde{u}$ and $\tilde{v}$ are in the same orbit of the action of G on $\tilde{\mathcal{MC}}(B)$. This defines the equivalence relation in equation 1.1.

We can verify immediately that if two deformation theories are governed by DG Lie algebras L_a and L_b , respectively, and if there is an (iso)morphism $L_a \longrightarrow L_b$ of DG Lie algebras, we get an (iso)morphism of deformation functors. But due to Kontsevich [2], there is a much stronger statement:

Theorem 1.1. *A quasi-isomorphism between two DG Lie algebras (in fact a quasi-isomorphism in the category of L_{∞} -algebras is sufficient) induces an isomorphism between the corresponding deformation functors.*

Below, to any associative algebra, we will assign two deformation functors Def_{rel} (deformations of relations) and Def_{ass} (associative deformations) that are governed

by DG Lie algebras L_{ass} and L_{rel} , respectively, and a deformation functor Def_{flat} , for which it is not clear, a priori, if it is governed by a DG Lie algebra. It was shown by Kontsevich [3] that the DG Lie algebras L_{rel} and L_{ass} are quasi-isomorphic, thus the deformation functors Def_{ass} and Def_{rel} are isomorphic. The aim of this paper is to show that the third one Def_{flat} is isomorphic to the both of them.

2 Deformations of the associative structure

Definition 2.1. A **deformation of the associative structure** of A with basis $B = (B, \mathfrak{m})$ is a B -linear map $\beta : A_B \otimes_B A_B \longrightarrow A_B$ with $\text{Im}(\beta) \subseteq A_{\mathfrak{m}}$ such that the sum $\alpha + \beta$ is associative as a product on A_B .

Definition 2.2. Two deformations β and β' of the associative structure of A with basis B are **equivalent**, if there exists a B -linear isomorphism $\Phi : (A_B, \alpha + \beta) \longrightarrow (A_B, \alpha + \beta')$ (in the unitary case, Φ has to be unitary) such that the following diagram commutes:

$$\begin{array}{ccc} & A & \\ \nearrow & & \nwarrow \\ A_B & \xrightarrow{\Phi} & A_B \end{array} \quad (*)$$

Maps without label are canonical.

It was discovered by Gerstenhaber, that the deformations of the associative structure of A are governed by a DG Lie algebra L_{ass} , namely the Hochschild cochain complex. The graded Lie bracket on this complex is now called the “Gerstenhaber bracket”. Let’s recall the modern definition of the Hochschild cochain complex:

In the sequel, M shall always denote the graded k -module $\downarrow A$ which is defined as $(\downarrow A)^{-1} = A$ and $(\downarrow A)^i = 0$, for $i \neq -1$. We have natural maps $\downarrow : A \longrightarrow M$ (degree -1) and $\uparrow : M \longrightarrow A$ (degree +1). For any graded k -module W , let $TW := \coprod_{n \geq 1} W^{\otimes n}$ be the (non-unitary) tensor algebra. On TW , we have a coproduct $\Delta : TW \longrightarrow TW \otimes TW$, defined by

$$w_1 \otimes \dots \otimes w_n \mapsto \sum_{i=1}^{n-1} (w_1 \otimes \dots \otimes w_i) \otimes (w_{i+1} \otimes \dots \otimes w_n).$$

Recall that the **bar construction** $\mathcal{B}A$ of A is the DG coalgebra (TM, Δ, Q) , where the degree one codifferential Q on TM is uniquely defined by its (co)restriction

$$\begin{aligned} Q_2 : M \otimes M &\longrightarrow M \\ m_1 \otimes m_2 &\mapsto -\downarrow \alpha(\uparrow m_1, \uparrow m_2). \end{aligned}$$

For more details, see [9]. The **Hochschild cochain complex** of A is the DG Lie algebra $L_{\text{ass}} := \text{Coder}(\mathcal{B}A) = \coprod_i \text{Hom}(M^{\otimes i}, M)$ of coderivations on the DG coalgebra

$\mathcal{B}A$. Its Lie bracket is the graded commutator and the (degree one) differential d_{ass} is given by $q \mapsto [Q, q]$. Observe that

$$L_{\text{ass}}^0(\mathfrak{m}) \cong \text{Hom}_{k\text{-Mod}}(A, A_{\mathfrak{m}}) \cong \{f \in \text{End}_{B\text{-Mod}}(A_B) \mid \text{Im } f \subseteq A_{\mathfrak{m}}\}.$$

Now, $L_{\text{ass}}^0(\mathfrak{m})$ is a nilpotent Lie subalgebra of $\text{End}_{B\text{-Mod}}(A_B)$. The Lie group G_{ass} associated to $L_{\text{ass}}(\mathfrak{m})^0$ is the subgroup

$$\{\phi \in \text{Aut}_{B\text{-Mod}}(A_B) \mid \phi = 1 + \psi \text{ for a } \psi \text{ with } \text{Im } \psi \subseteq A_{\mathfrak{m}}\}$$

of $\text{Aut}_{B\text{-Mod}}(A_B)$. The exponential map

$$\begin{aligned} \exp : L_{\text{ass}}(\mathfrak{m})^0 &\longrightarrow G_{\text{ass}} \\ f &\mapsto 1 + f + \frac{1}{2}f^2 + \dots \end{aligned}$$

is bijective. Its inverse is given by $1 + \psi \mapsto \ln(1 + \psi) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1} \psi^n}{n}$. Observe that

$$\begin{aligned} L_{\text{ass}}^1(\mathfrak{m}) &\cong \text{Hom}_{k\text{-Mod}}(A \otimes_k A, A_{\mathfrak{m}}) \\ &\cong \{\beta \in \text{Hom}_{B\text{-Mod}}(A_B \otimes_B A_B, A_B) \mid \text{Im } \beta \subseteq A_{\mathfrak{m}}\}. \end{aligned}$$

The group G_{ass} acts on $L_{\text{ass}}^1(B)$ by $\phi : \gamma \mapsto \phi \circ \gamma \circ (\phi^{-1} \otimes \phi^{-1})$. This is just the group action described more abstractly in Section 1. We identify deformations β and β' of α , if there exists an element ϕ of G_{ass} such that $\phi.(\alpha + \beta) = \alpha + \beta'$. (Be careful, in this case, we don't have $\beta' = \phi.\beta$!) This is just the equivalence relation in Definition 2.1. Verify that deformations of the associative structure of A with basis B correspond to Maurer-Cartan elements

$$\mathcal{MC}_{\text{ass}}(B) := \{\beta \in L_{\text{ass}}^1(\mathfrak{m}) \mid d_{\text{ass}}(\beta) + \frac{1}{2}[\beta, \beta] = 0\}.$$

Each element ϕ of G_{ass} , defines a bijection (that we shall also denote by ϕ)

$$\begin{aligned} \phi : \mathcal{MC}_{\text{ass}}(B) &\longrightarrow \mathcal{MC}_{\text{ass}}(B) \\ \beta &\mapsto \phi.(\alpha + \beta) - \alpha. \end{aligned}$$

Two elements β and β' in $\mathcal{MC}_{\text{ass}}(B)$ are equivalent, i.e. if $\beta' = \phi(\beta)$, for a $\phi \in G_{\text{ass}}$, if and only if they correspond to equivalent deformations of α . We define the deformation functor $\text{Def}_{\text{ass}} : \mathfrak{Art} \longrightarrow \mathfrak{set}$ as

$$\text{Def}_{\text{ass}}(B) := \mathcal{MC}_{\text{ass}}(B) / \sim.$$

3 Deformations of the relations

We can choose a free non-positively graded DGA resolution (R, s) of A , i.e. (R, s) is the free graded tensor algebra over a graded k -module together with a derivation s of degree $+1$ with $s^2 = 0$ such that (R, s) is acyclic in negative degrees and $H^0(R, s) \cong A$ via a fixed morphism $R^0 \longrightarrow A$. It can easily be verified that such a resolution exists and we will see later that there is even a canonical one.

Definition 3.1. A **deformation of the relations** of A with basis B is a degree one derivation δ of the free graded B -algebra $R_B = R \otimes_k B$ such that the following two conditions hold:

- (1) $\text{Im } \delta \in R \otimes_k \mathfrak{m}$,
- (2) $(s + \delta)^2 = 0$, i.e. the perturbed derivation $s + \delta$ on R_B is again a differential.

First, we want to prove that, for such a deformation δ , the complex $(R_B, s + \delta)$ is acyclic in negative degrees.

Proposition 3.2. *For each deformation δ of the relations of A with base B , the complex $(R_B, s + \delta)$ is a free resolution of $\tilde{A} := H^0(R_B, s + \delta)$ over B .*

Proof. The inclusions $\dots \subseteq \mathfrak{m}^2 \subseteq \mathfrak{m} \subseteq B$ give rise to a filtration of the complex $(R_B, s + \delta)$. The term $E_1^{p,q}$ of the spectral sequence assigned to this filtration is zero, for $p + q \neq 0$. The statement is thus a consequence of the Classical Convergence Theorem (see [10]) for spectral sequences. $\square$

Definition 3.3. Two Deformations δ_1 and δ_2 of the relations of A with basis B are **equivalent**, if there exist a B -linear isomorphism $\Phi : (R_B, s + \delta_1) \longrightarrow (R_B, s + \delta_2)$ of DG algebras such that the diagram

$$\begin{array}{ccc} & R & \\ \nearrow & & \nwarrow \\ R_B & \xrightarrow{\Phi} & R_B \end{array} \quad (**)$$

commutes.

Again, we will construct a DG Lie algebra, governing the deformations of the relations of A . Let L_{rel} be the graded module $\text{Der}(R) = \coprod_{i \in \mathbb{Z}} \text{Der}^i(R)$ of k -linear derivations of R . It carries a DGL structure, where the Lie bracket is again the graded commutator and the degree one differential d_{rel} is given by $\delta \mapsto [s, \delta]$. The complex $(L_{\text{rel}}, d_{\text{rel}})$ is called **(noncommutative) tangent complex**. As above, consider the DG Lie algebras $L_{\text{rel}}(B) := L_{\text{rel}} \otimes B$ and $L_{\text{rel}}(\mathfrak{m}) := L_{\text{rel}} \otimes \mathfrak{m}$. The Liegroup corresponding to the nilpotent Lie algebra $L_{\text{rel}}^0(\mathfrak{m})$ is given by

$$G_{\text{rel}} = \{\phi \in \text{Aut}_{B\text{-Alg}}(A_B) \mid \phi = 1 + \psi \text{ for a } \psi \text{ with } \text{Im } \psi \subseteq A_{\mathfrak{m}}\}.$$

Remark that in contrast to G_{ass} , the elements of G_{rel} are algebra homomorphisms. The exponential map

$$\begin{aligned} \exp : L_{\text{rel}}(\mathfrak{m})^0 &\longrightarrow G_{\text{rel}} \\ f &\mapsto 1 + f + \frac{1}{2}f^2 + \dots \end{aligned}$$

is bijective. The group G_{rel} acts on $L_{\text{rel}}^1(B)$ by $\phi : \gamma \mapsto \phi\gamma\phi^{-1}$. We identify deformations δ and δ' of the relations of A with basis B , if there exists an element of G_{rel} such that $\phi.(s + \delta) = s + \delta'$. Verify that the deformations of the relations of A are in one-to-one correspondence with the Maurer-Cartan elements:

$$\mathcal{MC}_{\text{rel}}(B) := \{\delta \in L_{\text{rel}}(\mathfrak{m})^1 \mid d_{\text{rel}}(\delta) + \frac{1}{2}[\delta, \delta] = 0\}$$

Each element ϕ of G_{rel} , defines a bijection (that we shall also denote by ϕ)

$$\begin{aligned} \phi : \mathcal{MC}_{\text{ass}}(B) &\longrightarrow \mathcal{MC}_{\text{ass}}(B) \\ \beta &\mapsto \phi.(\alpha + \beta) - \alpha. \end{aligned}$$

Two elements δ and δ' in $\mathcal{MC}_{\text{rel}}(B)$ are equivalent, i.e. if $\delta' = \phi(\delta)$, for a $\phi \in G_{\text{rel}}$, if and only if they correspond to equivalent deformations of the relations of A with basis B . We define the deformation functor $\text{Def}_{\text{rel}} : \mathfrak{Art} \longrightarrow \mathfrak{set}$ as

$$\text{Def}_{\text{rel}}(B) := \mathcal{MC}_{\text{rel}}(B) / \sim,$$

where $\sim$ denotes the relation generated by this group action. The following theorem, taken from [3], proves that the deformation functor does not depend on the chosen resolution R of A :

Theorem 3.4. *The tangent Lie algebras $L_{\text{rel},1}$ and $L_{\text{rel},2}$, corresponding to two different free DGA resolutions R_1 and R_2 of A are quasi-isomorphic as DGL algebras. As consequence, they define isomorphic deformation functors.*

4 Flat deformations

Definition 4.1. A **flat deformation** of A with basis B is a flat associative B -algebra $\tilde{A}$, together with a fixed isomorphism $\tilde{A} \otimes_B k \longrightarrow A$.

Example 4.2. For A commutative, the deformations of the affine scheme $X = \text{Spec } A$ over $\text{Spec } B$ are by definition cartesian diagrams

$$\begin{array}{ccc} \mathcal{X} & \longleftarrow & X \\ \downarrow g & & \downarrow \\ \text{Spec } B & \longleftarrow & * \end{array}$$

If $\mathcal{X} = \text{Spec } \tilde{A}$ is an affine flat scheme over $\text{Spec } B$, then $\tilde{A}$ is a flat deformation of A .

Definition 4.3. Two flat deformations $\tilde{A}_1$ and $\tilde{A}_2$ of A with basis B are **equivalent**, if there exists a B -algebra isomorphism $\Phi : \tilde{A}_1 \longrightarrow \tilde{A}_2$ such that the diagram

$$\begin{array}{ccc} & A & \\ \nearrow & & \nwarrow \\ \tilde{A}_1 & \xrightarrow{\Phi} & \tilde{A}_2 \end{array} \quad (***)$$

commutes.

Flat deformations of A are not directly related to a DG Lie algebra. We define the deformation functor $\text{Def}_{\text{flat}} : \mathfrak{Art} \longrightarrow \mathfrak{set}$ by

$$\text{Def}_{\text{flat}}(B) := \{\text{flat deformations of } A \text{ with basis } B\} / \sim .$$

4.1 From associative deformations to flat deformations

If β is a deformation of the associative structure of A with basis B , then $\tilde{A} := (A_B, \alpha + \beta)$ is a flat deformation of A . If β is equivalent to β' as deformation of α , then $(A_B, \alpha + \beta)$ is equivalent to $(A_B, \alpha + \beta')$ as flat deformation. Thus we have a natural transformation

$$F : \text{Def}_{\text{ass}} \longrightarrow \text{Def}_{\text{flat}}$$

of deformation functors. By construction, for any artinian basis B , the map $F(B) : \text{Def}_{\text{ass}}(B) \longrightarrow \text{Def}_{\text{flat}}(B)$ is injective.

4.2 From deformations of the relations to flat deformations

We fix a free DGA resolution (R, s) of A . This data contains an algebra map $\phi : R^0 \longrightarrow A$, inducing an isomorphism $H^0(R, s) \longrightarrow A$. Let δ be a deformation of the relations of A with basis B . We have seen that the DGA algebra $(R_B, s + \delta)$ is a free resolution of $\tilde{A} := H^0(R_B, s + \delta)$ over B .

Proposition 4.4. *The algebra $\tilde{A}$ is flat as B -module.*

Proof. For each B -module M and each $i > 0$, we have

$$\text{Tor}_i^B(\tilde{A}, M) = H^{-i}(R_B \otimes_B M) = 0,$$

which follows by the same spectral sequence argument as in the proof of Proposition 3.2. $\square$

The assignment $\delta \mapsto H^0(R_B, s + \delta)$ defines a natural transformation

$$G : \text{Def}_{\text{rel}} \longrightarrow \text{Def}_{\text{flat}}$$

of deformation functors. The next aim is to show that G is surjective. We will make use of Nakayama's Lemma for Artin algebras (that works without any finitely generated assumption). For the proof, see [5].

Lemma 4.5. *For a local artinian algebra $(B, \mathfrak{m})$ and any B -module, if $M \otimes_B B/\mathfrak{m} = 0$, then $M = 0$.*

In the sequel, for any B -module M , we will use the abbreviation $M' := M \otimes_B B/\mathfrak{m}$.

Lemma 4.6. *Consider a complex*

$$\cdots \longrightarrow M^{-2} \xrightarrow{d^{-2}} M^{-1} \xrightarrow{d^{-1}} M^0$$

of flat B -modules such that the induced complex (M', d') is acyclic (in degrees $\leq i$). Then, (M, d) is acyclic (in degrees $\leq i$ and $\text{Kern}(d^{i-1})$ is flat). The same statement is true, if we replace B by any local noetherian algebra under the additional assumption that each M^i is a finitely generated B -module.

Proof. First, we show that d^{-1} is surjective: We have a short exact sequence

$$0 \longrightarrow \text{Im}(d^{-1}) \longrightarrow M^0 \longrightarrow H^0(M) \longrightarrow 0.$$

Application of the right exact functor $\otimes_B B/\mathfrak{m}$ gives an exact sequence

$$\text{Im}(d^{-1})' \longrightarrow M'^0 \longrightarrow H^0(M)' \longrightarrow 0. \quad (4.2)$$

Since the epimorphism d'^{-1} factors through $\text{Im}(d^{-1})'$, the first arrow of the exact sequence 4.2 is surjective, hence $H^0(M)' = 0$. By Lemma 4.5, we get $H^0(M) = 0$. Next we show that $H^{-1}(M) = 0$: Since we have an exact sequence

$$0 \longrightarrow \text{Kern}(d^{-1}) \longrightarrow M^{-1} \longrightarrow M^0 \longrightarrow 0, \quad (4.3)$$

the flatness of M^0 and M^{-1} implies the flatness of $\text{Kern}(d^{-1})$. Applying the functor $\otimes_B B/\mathfrak{m}$ to the exact sequence 4.3, we get an exact sequence

$$0 \longrightarrow \text{Kern}(d^{-1})' \longrightarrow M'^{-1} \longrightarrow M'^0 \longrightarrow 0, \quad (4.4)$$

hence we have a natural isomorphism $\text{Kern}(d'^{-1}) \cong \text{Kern}(d^{-1})'$. Applying $\otimes_B B/\mathfrak{m}$ to the short exact sequence

$$0 \longrightarrow \text{Im}(d^{-2}) \longrightarrow \text{Kern}(d^{-1}) \longrightarrow H^{-1}(M) \longrightarrow 0,$$

we get an exact sequence

$$\text{Im}(d^{-2})' \longrightarrow \text{Kern}(d^{-1})' \longrightarrow H^{-1}(M)' \longrightarrow 0.$$

Since the epimorphism $M'^{-2} \longrightarrow \text{Kern}(d'^{-1})$ factors through $\text{Im}(d^{-2})'$, we see that the first map of the last exact sequence is surjective. Thus $H^{-1}(M)' = 0$. Again, by Lemma 4.5, $H^{-1}(M) = 0$. Continuing inductively in the same manner, we get that for each $i \leq 1$, the module $\text{Kern}(d^i)$ is flat over B and that $H^i(M) = 0$. $\square$

Proposition 4.7. *For each algebra B in $\mathfrak{Art}$, the map $G(B) : \text{Def}_{\text{rel}}(B) \longrightarrow \text{Def}_{\text{flat}}(B)$ is surjective.*

Proof. Let $f : B \longrightarrow \tilde{A}$ be a flat deformation of A with basis B . As graded algebra, set $\tilde{R} := R_B$. Remark that $\tilde{R}' = \tilde{R} \otimes_B k = R$.

We claim that we can find (1) a lifting $\tilde{\phi} : \tilde{R}^0 \longrightarrow \tilde{A}$ of ϕ , i.e. map of B -algebras such that $\tilde{\phi}' = \phi$, and (2) a differential $\tilde{s}$ on the graded algebra $\tilde{R}$ such that $\tilde{s}' = s$ and such that $\tilde{\phi}$ induces an isomorphism $H^0(\tilde{R}, \tilde{s}) \longrightarrow \tilde{A}$.

Since $\tilde{R}^0$ is a free B -algebra, a lifting $\tilde{\phi}$ of ϕ can be chosen. By Lemma 4.6, this lifting $\tilde{\phi}$ is surjective and thus, $\text{Kern } \tilde{\phi}$ is flat over B . It follows that $(\text{Kern } \tilde{\phi})' \cong \text{Kern } \phi$. We define the differential $\tilde{s}$ inductively, by choosing its values on the free generators x of R . For be a free generator x of degree -1 , there is a lifting $\tilde{r}$ of $s(x)$ in $\text{Kern } \tilde{\phi}$. Set $\tilde{s}(x) := \tilde{r}$. In this way, we define the component $\tilde{s}^{-1} : \tilde{R}^{-1} \longrightarrow \tilde{R}^0$. Applying Nakayama, as in the proof of Lemma 4.6, we see that $\text{Im}(\tilde{s}^{-1}) = \text{Kern } \tilde{\phi}$, thus we have a short exact sequence

$$0 \longrightarrow \text{Kern } \tilde{s}^{-1} \longrightarrow \tilde{R}^{-1} \longrightarrow \text{Kern } \tilde{\phi} \longrightarrow 0.$$

It follows that $\text{Kern } \tilde{s}^{-1}$ is flat over B and that $\text{Kern } s^{-1} = (\text{Kern } \tilde{s}^{-1})'$. Now, for free generators x of R of degree -2 , we can define $\tilde{s}^{-2}(x)$ as a lift of $s^{-2}(x)$ in $\text{Kern } \tilde{s}^{-1}$ and so on. Inductively, for each $n > 0$, we can define the values $\tilde{s}^{-n}(x)$, for free generators of R of degree $-n$ in such a way that $\text{Kern } \tilde{s}^{-n}$ is flat over B and such that the complex

$$\text{Kern } \tilde{s}^{-n} \longrightarrow \tilde{R}^{-n} \longrightarrow \dots \longrightarrow \tilde{R}^0$$

is a resolution of $\tilde{A}$. Thus the differential $\tilde{s}$ can be constructed as desired.

Finally, we see that $\tilde{s}$ equals $s + \delta$, for a deformation δ of the relations of A with basis B . $\square$

5 The canonical free DGA resolution

Consider a differential graded coalgebra C with comultiplication Δ and degree one codifferential q . Set W to be the shifted module $C[-1] = \uparrow C$. Recall that the **cobar construction** ΩC of C is the free graded tensor algebra TW together with the degree one differential d , which is uniquely defined by its corestrictions $d_1 := \uparrow q \downarrow : W \longrightarrow W$ and $d_2 := \uparrow^{\otimes 2} \Delta \downarrow$. The cobar construction is a functor


$$\Omega : \text{DG-Alg} \longrightarrow \text{DG-Coalg}.$$

The following statement can be found in [4] or [9].

Theorem 5.1. *(1) Bar and cobar construction form an adjoint pair $(\Omega, \mathcal{B})$ of functors. (With appropriate model structures on the categories DG-Alg and DG-Coalg they even form a Quillen-functor.)*

- (2) For any DG algebra A , the natural map $\Omega\mathcal{B}A \longrightarrow A$ is a quasi-isomorphism. I.e. $\Omega\mathcal{B}A$ is a canonical free DGA resolution of A .

It is worth, studying more closely the canonical free DGA resolution $\Omega\mathcal{B}A$ of an associative k -algebra A . As earlier, set $M := \downarrow A$. Recall that the bar construction $\mathcal{B}A$ is the DG coalgebra TM with coalgebra structure as in Section 2. As graded object, the complex $\Omega\mathcal{B}A$ equals $T \uparrow (TM)$. Picturally:



For $i \leq 0$, the degree i component $\Omega\mathcal{B}A^i$ of $\Omega\mathcal{B}A$ is the sum over all terms of the form $\uparrow (M \otimes \dots \otimes M) \otimes \dots \otimes \uparrow (M \otimes \dots \otimes M)$, containing $n = 1, 2, 3, \dots$ copies of M and $n + i$ arrows. We call the number n the **polynomial degree**. The polynomial degree is a second grading of the algebra $\Omega\mathcal{B}A$. A third one is given by the number of arrows arising in a summand. The three gradings are related by

$$\text{degree} = \text{number of arrows} - \text{polynomial degree}.$$

The differential d of $\Omega\mathcal{B}A$ has two components. One component d_{sw} of polynomial degree -1 and one component d_s increasing the number of arrows by one. Since $\Omega\mathcal{B}A$ is the free tensor algebra over $\uparrow (TM)$, the differential is uniquely defined by its values on terms of the form $\uparrow (M \otimes \dots \otimes M)$ (upper diagonal). We have

$$d_s(\uparrow (m_1 \otimes \dots \otimes m_n)) = \sum_{j=1}^{n-1} (-1)^j \uparrow (m_1 \otimes \dots \otimes m_j) \otimes \uparrow (m_{j+1} \otimes \dots \otimes m_n)$$

and d_{sw} is just the shifted bar complex, i.e.

$$d_{sw}(\uparrow (m_1 \otimes \dots \otimes m_n)) = \sum_{j=1}^{n-1} (-1)^{j-1} \uparrow (m_1 \otimes \dots \otimes m_{j-1} \otimes m_j \cdot m_{j+1} \otimes \dots \otimes m_n),$$

where $m_j \cdot m_{j+1}$ stands for $\downarrow (\uparrow m_j \cdot \uparrow m_{j+1})$. The natural algebra map $p : \Omega\mathcal{B}A \longrightarrow A$ given by the adjoint functor property of the bar and cobar construction is the map from the degree zero component to A given by

$$p : \uparrow m_1 \otimes \dots \otimes \uparrow m_n \mapsto \uparrow m_1 \cdot \dots \cdot \uparrow m_n.$$

The complex $\Omega\mathcal{B}A$ is filtered by the polynomial degree and the associated differential graded ring is the sum over the rows in the above picture. All rows except the first one are acyclic split. The splitting is given by the formular (see Chapter 3.4 of [9] or Page 36 of [4])

$$\begin{aligned} & \uparrow (m_{11} \otimes \dots \otimes m_{1i_1}) \otimes \dots \otimes \uparrow (m_{k1} \otimes \dots \otimes m_{ki_k}) \mapsto \\ & \left\{ \begin{array}{ll} 0 & \text{for } i_1 \neq 1 \\ \uparrow (m_{11} \otimes m_{21} \otimes \dots \otimes m_{2i_2}) \otimes \dots \otimes \uparrow (m_{k1} \otimes \dots \otimes m_{ki_k}) & \text{for } i_1 = 1 \end{array} \right. \end{aligned}$$

Again by convergence of the spectral sequence obtained from the polynomial filtration, the map p is a quasi-isomorphism. By the same reason, the natural inclusion $A = \uparrow M \longrightarrow \Omega\mathcal{B}A$ is also a quasi-isomorphism. Of course it is not an algebra map. The proof of the following theorem is encrypted in [3]:

Theorem 5.2. *There is a natural quasi-isomorphism*

$$L_{\text{rel}} = \text{Der}(\Omega\mathcal{B}A) \longrightarrow \text{Coder}(\mathcal{B}A) = L_{\text{ass}}$$

respecting the DGL-structures. In consequence, we get an isomorphism $\text{Def}_{\text{rel}} \longrightarrow \text{Def}_{\text{ass}}$ of deformation functors.

Proof. The quasi-isomorphism $p : \Omega\mathcal{B}A \longrightarrow A$ induces a quasi-isomorphism

$$\text{Der}(\Omega\mathcal{B}A) = \text{Hom}(\uparrow \mathcal{B}A, \Omega\mathcal{B}A) \longrightarrow \text{Hom}(\uparrow \mathcal{B}A, A) \cong \text{Hom}(\mathcal{B}A, \downarrow A) = \text{Coder}(\mathcal{B}A).$$

Check that it respects the DGL structures. $\square$

In the sequel, we will form the set $\mathcal{MC}_{\text{rel}}(B)$ with respect to the canonical free DGA resolution $\Omega\mathcal{B}A$ of A . By Theorem 1.1, the quasi-isomorphism of Theorem 5.2 induces, for each $B \in \mathfrak{Art}$, a bijection $E : \text{Def}_{\text{rel}}(B) \longrightarrow \text{Def}_{\text{ass}}(B)$. The knowledge of the structure of $\Omega\mathcal{B}A$ allows us to describe easily the inverse E^{-1} : Let $\beta \in \mathcal{MC}_{\text{ass}}(B)$ be a Maurer-Cartan element. The differentials of $\Omega\mathcal{B}(A_B, \alpha)$ and $\Omega\mathcal{B}(A_B, \alpha + \beta)$ on the graded algebra $\Omega\mathcal{B}A_B$ only differ by a component $\delta(\beta)$ in south-west direction. We have $\delta(\beta) \in \mathcal{MC}_{\text{rel}}(B)$ and the quasi-isomorphism of Theorem 5.2 maps the class of $\delta(\beta)$ to β . Thus, E maps the class $[\delta(\beta)]$ of $\delta(\beta)$ in $\text{Def}_{\text{rel}}(B)$ to the class $[\beta]$ of β in $\text{Def}_{\text{ass}}(B)$.

Now we are able to prove the main statement:

Theorem 5.3. *For any local commutative artinian algebra B , the diagram*

$$\begin{array}{ccc} \text{Def}_{\text{rel}}(B) & \xrightarrow{E} & \text{Def}_{\text{ass}}(B) \\ & \searrow G & \downarrow F \\ & & \text{Def}_{\text{flat}}(B) \end{array}$$

commutes and all the arrows are bijections.

Proof. For $[\beta] \in \text{Def}_{\text{ass}}(B)$, we have $G(E^{-1}([\beta])) = G([\delta(\beta)]) = H^0(\Omega\mathcal{B}A_B, d + \delta(\beta)) \cong (A_B, \alpha + \beta) \cong F([\beta])$. This proves the commutativity. Since E is an bijective, G is surjective and F is injective, we see that F and G are also isomorphisms. $\square$

Corollary 5.4. *The three deformation functors Def_{ass} , Def_{rel} and Def_{flat} are isomorphic.*

By Theorem 5.3, the functors E , F and G are in particular smooth. Thus, by an observation of Schlessinger [8], Corollary 5.4 is still true, if we replace the three deformation functors by their extensions to certain categories of projective limits of Artin algebras, for example the category of complete noetherian local algebras with residue field k .

References

- [1] M. Gerstenhaber; S.D. Schack: *Algebraic cohomology and deformation theory*, Michiel Hazewinkel and Murray Gerstenhaber (ed.), Deformation Theory of Algebras and Structures and Applications, Kluwer (1988), 11–264.
- [2] Maxim Kontsevich: *Deformation quantization of Poisson manifolds I*, preprint q-alg/9709040.
- [3] Maxim Kontsevich: *Topics in algebra - deformation theory*, notes by A. Weinstein, <http://wuarchive.wustl.edu/doc/techreports/mcgill.ca/math/fox/TFOX.HTM>.
- [4] Kenji Levêfre-Hasegawa: *Sur les A_∞ -catégories*, Thèse Université Paris 7, (2003).
- [5] Marco Manetti: *Deformation of singularities via differential graded Lie algebras*, notes (2001).
- [6] Frank Schuhmacher: *Deformations of L_∞ -algebras*, preprint
- [7] Frank Schuhmacher: *Deformation of singularities via L_∞ -algebras*, preprint
- [8] Michael Schlessinger: *Functors of Artin Rings*, Trans. Amer. Math. Soc. 130 (1968) 208-222.
- [9] Vladimir Alekseevich Smirnov: *Simplicial and operad methods in algebraic topology*, Translation of Mathematical Monographs, vol. 198, AMS (2001).
- [10] Charles A. Weibel: *An introduction to homological algebra*, Cambridge studies in advanced mathematics 38, (1994).